UNCLASSIFIED
NSN 7540-01-280-5500

OR REPORT

protein engineering

17. SECURITY CLASSIFICATION

19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED

20. LIMITATION OF ABSTRACT

UL

Standard Form 298 (Rev. 2-89) Prescribed by ANSI Std. 239-18

SECURITY CLASSIFICATION

UNCLASSIFIED

OF THIS PAGE

MEMORANDUM TRANSMITTAL

U. S. Army Research Office ATTN: AMXRO-ICA (Hall) P. O. Box 12211 Research Traingle Park, NC 27709-2211

Reprint (Orig + 2 copies)	Technical Report (Orig + 2 copies)
Manuscript (1 copy)	Final Progress Report (Orig + 2 copies
	Other (1 copy)
CONTRACT/GRANT NUMBER:	DAAG5597-0173
REPORT TITLE: Molecular Geneti	c Approaches to Biomolecular Materials
is forwarded for your information.	
SUBMITTED FOR PUBLICATION TO	(applicable only if report is manuscript):

Hagan Bayley, Ph.D. Professor

MOLECULAR GENETIC APPROACHES TO BIOMOLECULAR MATERIALS

FINAL PROGRESS REPORT

HAGAN BAYLEY

NOVEMBER 15, 2000

DAAG5597-0173

THE TEXAS A&M UNIVERSITY SYSTEM HEALTH SCIENCE CENTER

APPROVED FOR PUBLIC RELEASE;
DISTRIBUTION UNLIMITED.

THE VIEWS, OPINIONS, AND/OR FINDINGS CONTAINED IN THIS REPORT ARE THOSE OF THE AUTHOR(S) AND SHOULD NOT BE CONSTRUED AS AN OFFICIAL DEPARTMENT OF THE ARMY POSITION, POLICY, OR DECISION, UNLESS SO DESIGNATED BY OTHER DOCUMENTATION.

FINAL REPORT

GRANT #: DAAG5597-0173

<u>PRINCIPAL</u> <u>INVESTIGATOR</u>: Hagan Bayley, PhD; AASERT student: George Miles

<u>INSTITUTION</u>: The Texas A&M University System Health Science Center

GRANT TITLE: The student was engaged in research under the ARO grant DAAG5597-0173: Molecular genetical approaches to biomolecular materials

AWARD PERIOD: June 1 1997 - May 31 2000

OBJECTIVE: The leukocidins are bicomponent pore-forming toxins that are structurally related to staphylococcal alphahemolysin, a single component toxin that has been studied in detail by the Bayley group. alpha-Hemolysin has many applications in the area of biomolecular materials. Therefore, leukocidins have been characterized to expand the set of tools available for materials engineering.

APPROACH: Staphylococcal LukF and LukS together form transmembrane pores in lipid bilayers. The approach taken in the AASERT project was to obtain genes for the two components of the toxin, to express them in E. coli, to characterize the heterologously expressed pores by bilayer recording, and to use molecular genetic technology to explore structure-function relationships in these proteins.

ACCOMPLISHMENTS: The first three of the four objectives have been accomplished and have written up as a paper for Biochemistry. The genes for LukF and LukS were amplified from Staphylococcus aureus genomic DNA, placed in expression vectors and sequenced. The genes were also constructed by using synthetic oligonucleotides as building blocks. latter was done so that the sequence homology at the nucleotide level could be increased, while maintaining the original amino acid sequences. This will allow gene shuffling experiments to be performed. The LukF and LukS genes were expressed in E. coli cells or by in vitro translation in an E. coli S30 system. The products were purified either native or His-tagged form. The electrical properties various combinations of subunits were explored by singlechannel recording. The LukF/ LukS pore is larger than that formed by alpha-hemolysin and has other distinct properties including ion selectivity and gating characteristics.

CONCLUSIONS & SIGNIFICANCE: The availability of milligram amounts of the leukocidins will be useful for membrane protein engineering studies that will increase the diversity

of pores that can be formed by alpha-hemolysin and its relatives.

PATENT INFORMATION: not applicable

, , ·

AWARD INFORMATION: none

PUBLICATIONS:

Miles, G., Braha, O., Cheley, S. and Bayley, H. (2000) Biochemical and biophysical characterization of recombinant LukF/ LukS, a bicomponent pore-forming toxin, Biochemistry, submitted